

Spectral line and continuum radiation propagation in a clumpy medium

John Conway

Onsala Space Observatory, Sweden

Moshe Elitzur

Department of Physics and Astronomy, University of Kentucky, USA

Rodrigo Parra

Onsala Space Observatory, Sweden

April 15, 2004

Abstract. We discuss the propagation of spectral line and continuum radiation in a clumpy medium and give general expressions for the observed absorption or emission from a cloud population. We show that the affect of the medium clumpiness can usually be characterised by a single number multiplying the mean column opacity. Our result provides a simpler proof and generalization of the result of Martin et al. (1984). The formalism provides a simple way to understand the effects of clumping on molecular line profiles and ratios; for example how clumping effects the interpretation of $^{13}\text{CO}(1-0)$ to $^{12}\text{CO}(1-0)$ line ratios. It also can be used as a propagation operator in physical models of clumpy media where the incident radiation effects the spectral line emissivity. We are working to extend the formalism to the propagation of masers in a clumpy medium, but in this case there are special difficulties because formal expectation values are not characteristic of observations because they are biased by rare events.

Keywords: molecular lines, spectra

1. Introduction

The problem of the propagation of radiation in a non-uniform medium is a general one in astrophysics, which arises in many different contexts. We consider here the common case that the medium consists of clumps or clouds with a low volume filling factor, but potentially high covering factor. These clumps or clouds may possess a range of properties. This paper gives a summary of our derivation and results, a more detailed proof will be published elsewhere (Conway, Elitzur and Parra 2004, in prep).

The specific problem we address is the expected response of a telescope beam observing a low filling factor clumpy region where the beamwidth is much larger than the typical cloud diameter. In this case we can show that that expectation value of the beam average equals the ensemble expectation value over many realisations of observations



© 2008 Kluwer Academic Publishers. Printed in the Netherlands.

along the line of sight (LOS). We first (see Section 2) analyse the case of continuum (e.g. dust) clouds absorbing background radiation, then spectral line absorption (Section 3). In Section 4 we consider spectral line emission and apply the formalism to the interpretation of CO line ratios. Finally in Section 5 we draw conclusions and describe future work.

2. Continuum Absorption Radiation Transfer

Let us consider continuum radiation transfer in a clumpy medium of dust clouds. First assume that all clouds/clumps are identical. Assume further that radiation I entering a clump emerges as $I \cdot R$, where the clump response is R . In terms of cloud opacity $R = e^{-\tau}$. Hitting successively k clumps that all have the same response factor produces the aggregate response R^k . Implicit in this assumption is that the interaction with the radiation does not change the clump response factor R . It can be shown that provided the clump volume filling factor is small the probability of getting k clumps along a given line of sight always obeys a Poisson distribution, characterised by one number, the mean $\bar{N}$ along a LOS. In this case of one clump type it can be shown that the mean response is

$$\bar{R} = e^{-t}, \quad t = \bar{N}(1 - e^{-\tau}) \quad (1)$$

as previously derived by Natta & Panagia (1992).

Next consider the case of many types of clumps with response factors R_i (these could be clump types having different optical depths) each obeying Poisson statistics. If the fraction of clumps of each type is f_i then (see Conway, Elitzur and Parra 2004, in prep)

$$\bar{R} = \exp(\bar{N}[R_{\text{av}} - 1]), \quad \text{where} \quad R_{\text{av}} = \sum_i f_i R_i \quad (2)$$

This means that the problem of a mixture of cloud optical depths (τ_i) is equivalent to that of the single cloud type having an average τ_{av} obtained from

$$R_{\text{av}} = e^{-\tau_{\text{av}}} = \sum_i f_i e^{-\tau_i} \quad (3)$$

We can generalize the result in Eq. (2) to the case where the clump type varies continuously. For instance if the clump type is described by continuous variables t and r (which may correspond for instance to peak optical depth and distance from the centre of clouds which have spatial structure) then we have

$$R_{\text{av}} = \int \int R(r, t) \eta(r, t) dr dt = \int \int e^{-\tau(r, t)} \eta(r, t) dr dt \quad (4)$$

and likewise for any number of dimensions of continuous parameters, where η is the distribution function of clump types.

3. Line Radiation Transfer - Absorption

In the spectral line absorption problem we aim to calculate the expectation value of a frequency dependent absorption response function given a population of clouds. This cloud population has in general a distribution of central LOS velocities and a range of spectral profiles and spatial structures. Observed absorption lines are generally published as a function of Doppler velocity v , the response at velocity $R(v)$ being understood to represent the response at observing frequency $\nu = \nu_0(1 + v/c)$ where ν_0 is the line's rest frequency. Let us concentrate on calculating the response at a particular velocity v , $R(v)$. In principle clouds having all possible centre velocities v' contribute to the spectral response at velocity v . The centre velocity of each cloud v' can simply be thought of as yet another continuous variable describing the cloud properties, so the full distribution function becomes $\eta(r, t, v')$. For spectral line clouds the response function is $R_v(r, t, v')$ which if cloud types are defined to have fixed spectral profiles which don't vary with centre velocity can be written $R(r, t, v - v')$. Hence applying Eq. (2) and (4) we obtain

$$\bar{R}(v) = \exp(\bar{N}[R_{\text{av}}(v) - 1]) \quad (5)$$

$$\text{where } R_{\text{av}}(v) = \int \int \int R(r, t, v - v') \eta(r, t, v') dv' dr dt \quad (6)$$

As before $\bar{N}$ is the mean number of clouds (at any velocity) per LOS. $R_{\text{av}}(v)$ can be interpreted at each r, t value as a convolution of the cloud velocity profile for this r, t , $R(r, t, v)$ with the velocity distribution function of the cloud population $\eta(r, t, v)$ followed by an integration over all r, t (and any other continuous cloud parameters).

If the fractional distribution of clouds of different kinds is independent of velocity, then $\eta(r, t, v) = f(r, t)q(v)$ is a separable function. In this case we can define a mean cloud spectral response

$$R_s(v) = \int \int R(r, t, v) f(r, t) dr dt \quad (7)$$

In this separable case the expectation value of the emerging spectrum is

$$\bar{R}(v) = \exp(-\lambda(v) * C(v)) \quad (8)$$

In this expression $*$ denotes convolution, $\lambda(v) = \bar{N}q(v)$ describes the cloud velocity distribution; more precisely defined as the mean number of clouds per unit velocity per LOS. The quantity $C(v) = 1 - R_s(v) = \langle 1 - e^{-\tau(v)} \rangle$, where $\tau(v)$ is the opacity profile of each clump type, and the brackets denote an average over the clump population. The $C(v)$ can be interpreted as the mean cloud emission profile. In the normal case that the $\lambda(v)$ distribution is much broader in velocity than $C(v)$ we obtain the good approximation that $\bar{R}(v) \approx \exp(-C_A \lambda(v))$ where $C_A = \int C(v) dv$ has dimensions of velocity and can be considered the equivalent width of the mean cloud profile.

Another way to view the effect of clumping is to note that

$$\bar{R}(v) \approx \exp(-K \bar{\tau}_{tot}(v)) \quad (9)$$

where $\bar{\tau}_{tot}(v) = \lambda(v) * \langle \tau(v) \rangle$ is the mean line of sight opacity and K is a factor taking into account clumping which is

$$K = \frac{\int \langle 1 - e^{-\tau(v)} \rangle dv}{\int \langle \tau(v) \rangle dv} \quad (10)$$

Remarkably we find that the effect of an arbitrarily complex clump distribution can usually be reduced to a single number, K . This factor K reduces the effective opacity of a clumpy medium compared to a smooth gas of the same mean column density. Martin et al. (1984) partially derived our results, but their derivation was less direct and the full generality of the result was not stated.

4. Emission Line Spectra

The expressions for opacity derived above can be adapted (Conway, Elitzur and Parra 2004, in prep) to predict the emissivity in the case of a population of clouds with a mixture of excitation temperatures and statistical properties along the line of sight. Here we discuss only the simplest case that all clouds have the same excitation temperature T_e . In this case the observed spectral brightness temperature is

$$T(v) = T_e[1 - \bar{R}(v)] = T_e[1 - \exp(-K \bar{\tau}_{tot}(v))] \quad (11)$$

To illustrate the impact of clumping on observed line profiles and line ratios we show an example in Fig 1 involving the $^{12}\text{CO}(1-0)$ and $^{13}\text{CO}(1-0)$ lines. In this example the cloud population consists of only one clump

type with a Gaussian opacity velocity profile $\tau(v)$, and a much broader Gaussian $\lambda(v)$ cloud velocity distribution. The expectation value of the total opacity along each LOS $\bar{\tau}_{tot}(v) = \lambda(v) * \tau(v)$ is therefore also a Gaussian whose peak we term the 'peak total opacity'.

The different panels for Fig 1 have different combinations of the $^{12}\text{CO}(1-0)$ peak total and peak cloud opacities. The ratio of intrinsic $^{12}\text{CO}(1-0)$ and $^{13}\text{CO}(1-0)$ opacities is set to 60 so that in all the panels of Fig 1, the $^{13}\text{CO}(1-0)$ total and cloud opacities are $\ll 1$ and hence are optically thin. In all cases therefore the $^{13}\text{CO}(1-0)$ line profiles are Gaussian with peak value (in units of T_e) equal to the total $^{12}\text{CO}(1-0)$ opacity divided by 60. In contrast the $^{12}\text{CO}(1-0)$ line profiles and their ratio to the $^{13}\text{CO}(1-0)$ lines depend critically on the clump opacity which affects K . Only for the case in the top left panel, which has both a low total and low clump opacity, is the observed line ratio equal to the intrinsic value. If either or both total and clump opacities are larger than 1 then a smaller line ratio is observed, but it is impossible just from the line ratio at line centre to distinguish the different cases.

An important consequence of Eq. (9) is that when the cloud velocity distribution is much broader than individual clouds it is in principle impossible from the $^{12}\text{CO}(1-0)$ and $^{13}\text{CO}(1-0)$ spectra alone to separate the effects of clumping and variations in the intrinsic $^{12}\text{CO}(1-0)/^{13}\text{CO}(1-0)$ opacity ratio. In going from panels on the top row with low opacity clumps to the bottom row with large clumps the $^{12}\text{CO}(1-0)$ clump factor K is reduced. The resulting spectra are the same as if there was no clumping but the intrinsic $^{12}\text{CO}(1-0)/^{13}\text{CO}(1-0)$ opacity ratio was reduced.

5. Conclusions and Future Work

We have derived general expressions for both continuum and spectral line radiation propagation in a clumpy medium. These expressions hold provided only that the clumps have a small volume filling factor and that the local emissivity is not itself affected by the local line spectra. Eq. (8), (9) and (10) provide simple expressions that can be used by observers to interpret observed line profiles. In Section 4 we illustrated the application of this equation to CO line emission profiles from a clumpy medium. Although the results presented in Section 4 are not new our equations provide a way to understand line profiles from clumpy media in a systematic and quantitative way.

We are working to extend our formalism to predict variances and higher order statistics of spectra from clumpy media. We also hope to use our formalism as a propagator term in physical spectral line models




Figure 1. Example $^{12}\text{CO}(1-0)$ and $^{13}\text{CO}(1-0)$ line profiles (shown respectively by solid and dashed lines) for different combinations of cloud and mean column opacities. To make them more visible the $^{13}\text{CO}(1-0)$ profiles have been multiplied by 10. The intrinsic ratio of opacity in the two transitions is assumed to be 60. Both the internal cloud opacity profiles of the clouds in the two transitions and the cloud velocity distribution are assumed to be Gaussian with the latter having a FWHM 10 times larger. The top row shows the case of low opacity in each $^{12}\text{CO}(1-0)$ cloud, and a varying mean *total* $^{12}\text{CO}(1-0)$ opacity. The bottom row shows the case of high opacity $^{12}\text{CO}(1-0)$ clouds. Note the different vertical scales of each plot. Only when the total and cloud $^{12}\text{CO}(1-0)$ opacities are both less than one is the observed $^{12}\text{CO}(1-0)$ to $^{13}\text{CO}(1-0)$ ratio close to the intrinsic value of 60.

of clumpy media. In general in such models emissivity will depend on the local spectrum so such models must be iterative. Finally we are actively working (see Parra et al. these proceedings) on the problem of maser propagation in a clumpy medium. Formally exactly the same equations apply as in the thermal case, the only difference is the use of negative opacities. However as well as physical complications of maser saturation there is a more subtle effect that even for unsaturated masers the formal means predicted by the equations are not good predictors of observations. The reason for this is that these mean values are dominated by events causing very large amplification which are also very improbable, such that they never occur within a typical source.

References

- Martin H.M., Hills R.E., & Sanders D.B., 1984, MNRAS, 208, 35
 Natta A. & Panagia N., 1984, ApJ, 287, 228